

Spatial distribution analysis of the TROPOMI Aerosol Layer Height: A pixel-by-pixel comparison to EARLINET and CALIOP observations

K. Michailidis¹, M.-E. Koukouli¹, D. Balis D.¹, M. de Graaf², J.P. Veefkind J. P.²

¹Laboratory of Atmospheric Physics, Physics Department, AUTH, Greece

²Royal Netherlands Meterological Institute (KNMI), De Bilt, the Netherlands

O4. Synergistic use of multiple instruments and techniques, networks and campaigns
 27-Jun, 14:15
 Monday 04 P07

Instrumentation and Datasets




Scope of study: Validation of TROPOMI Aerosol layer height (ALH) product with the retrievals of the GB lidar systems of EARLINET (+ CALIPSO)

* Lidar profiles from the EARLINET & CALIPSO mission are a good source of data for validating retrieved ALHs from TROPOMI S5-P



Vertical structure of the atmosphere

* Offer high accuracy and validation results **BUT** their geographical coverage is spatial limited.



Study Period: April 2018 - Sep 2021


- The Lidar measurements used for the validation are analyzed centrally by EARLINET's SCC algorithm, following common QA procedures before being upload on EARLINET DB.
- Only data labeled as Level-2 are included in the validation process.

Only CALIOP dust plume heights over central-easten Mediterranean are analyzed, filter out by Feature_Classification_Flags.



A total of 8 coincident CALIOP lidar measurements are presented.

(*validation procedure following Nanda et al., 2020)



The 30th International Laser Radar Conference (ILRC) virtual conference, June 26th – July 1st, 2022.

Validation Methodology

EARLINET

















GOME-2



Optionally

AERONET



HYSPLIT

Validation strategy and development

TROPOMI

- EARLINET DB files up to September 2021 are analyzed here
- New lidar files have been just recently uploaded to the SCC & EARLINET DB by stations
 - We are currently collecting and filtering the new lidar data

Quality assured lidar data - Single Calculus Chain (SCC)

- We use the backscatter profiles (1064 or 532nm) to retrieve aerosol layer height.
 - High-quality aerosol optical products (Level-2)

TROPOMI pixels QA>=0.5 & additional flags

- Sunglint effect, snow/ice pixels, Cloud screening


Co-location criteria:

- Spatially averaged TROPOMI pixels in a radius of 150km
- A maximum time difference of ±4h for collocation pairs

Lidar weighted height


$$ALH_{bsc} = \frac{\int\limits_{z_{i=1}}^{z=n} z_{i} \cdot \beta_{aer,i}(z) dz}{\int\limits_{z_{i=1}}^{z=n} \beta_{aer,i}(z) dz}$$





Validation Results






Study Period: April 2018 - Sep 2021

34 collocated cases

TROPOMI pixels	R	Bias (km)
Sea	0.9	-1.017
Land & Sea	0.6	-1.459



- * EARLINET & CALIOP calculated height is consistently lower than TROPOMI ALH.
- * Different instrument sensitivity lidars against TROPOMI/S5P.
- * Over land, TROPOMI ALH becomes unreliable for increasing surface albedo.

8 day/cases - 521 collocated pixels

The 30th International Laser Radar Conference (ILRC) virtual conference, June 26th – July 1st, 2022.

Results - A case study: Dust event over Greece (22 June 2021)






PollyXT 24/7 lidar system
PANGEA station




National Observatory of Athens - NOA Location: [23.31°E / 35.86°N], Antikythera, Greece Date: 2021-06-22T11:31:00Z-2021-06-22T13:01:00Z




VIIRS/Suomi
True color image



TROPOMI ALH



TROPOMI UVAI



CALIPSO overpass over Greece 2021-06-22 12:21:19Z - 12:34:48


Aerosol Height retrievals:

TROPOMI: 2.55±0.4km

PollyXT: 3.01km

CALIPSO: 3.98km


Under a homogeneous cloud-free the mean TROPOMI ALH agree well with lidar ALH



Summary and conclusions



<u>Highlight:</u> A smoke case - Greek fire during summer 2021 (Aug 8)



- * The dispersion of the smoke plume is clearly visible.
- * High UVAI values (>5.0)
- * ALH values ranging: 1.5 2.5km.
- * No cloud detected during this event.



- ✓ The TROPOMI EARLINET study is focused on selected desert dust cases and fires plumes.
- ✓ EARLINET & CALIOP calculated aerosol height is consistently lower than TROPOMI ALH.
- ✓ Several cases between Oct 2021 May 2022 have been characterized as Saharan dust cases and are expected to be analysed and compared to measurements by EARLINET stations.
- ✓ We have investigated the optimum search radius around an EARLINET stations for TROPOMI, considering the high spatial resolution of instrument.